arXiv:1411.5505v1 [math.AP] 20 Nov 2014

SELF-SIMILAR BLOW-UP SOLUTIONS OF THE KPZ EQUATION

ALEXANDER GLADKOV

Abstract. In this paper we consider self-similar blow-up solutions for the generalized deterministic KPZ equation $u_t = u_{xx} + \lambda|u_x|^q, \lambda > 0, q > 2$. The asymptotic behavior of self-similar solutions are studied.

1. Introduction

We consider the generalized deterministic KPZ equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \lambda \left|\frac{\partial u}{\partial x}\right|^q, \quad (x,t) \in S_T := \mathbb{R} \times (0,T), \tag{1.1}$$

where λ, q and T are real constants, $\lambda > 0$, $q > 2$, $T > 0$. Equation (1.1) was first considered in the case $q = 2$ by Kardar, Parisi and Zhang [12] in connection with the study of the growth of surfaces. When $q = 2$ (1.1) has since been referred to as the deterministic KPZ equation. For $q \neq 2$ it also called the generalized deterministic KPZ equation or Krug-Spohn equation because it was introduced in [13]. We refer to review article [8] for references and a detailed historical account of the KPZ equation.

The existence and uniqueness of a classical solution of the Cauchy problem for (1.1) with $q = 1$ and initial function $u_0 \in C_0^3(\mathbb{R}^n)$ was proven in [4]. This result was extended to $u_0 \in C^2(\mathbb{R}^n) \cap W^{2,\infty}(\mathbb{R}^n)$ and $q \geq 1$ in [1] and to $u_0 \in C(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ and $q \geq 0$ in [9]. Several papers [2], [3], [5], [6], [7] were devoted to the investigation of the Cauchy problem for irregular initial data, namely for $u_0 \in L^p(\mathbb{R}^n)$, $1 \leq p < \infty$ or for bounded measures. The existence and uniqueness of a solution to the Cauchy problem with unbounded initial datum are proved in [10]. To confirm the optimality of obtained existence conditions the authors of [10] analyze the asymptotic behavior of self-similar blow-up solutions of (1.1) for $q < 2$.

In this paper for $q > 2$ we investigate the asymptotic behavior of self-similar blow-up solutions of (1.1) of the following form

$$u(x,t) = (T_0 - t)^\alpha f(\xi), \quad \text{where } \xi = |x|(T_0 - t)^\beta. \tag{1.2}$$

Here T_0 is any positive constant. After substitution (1.2) into (1.1) we find that

$$\alpha = \frac{q-2}{2(q-1)}, \quad \beta = -\frac{1}{2} \tag{1.3}$$

and f should satisfy the following equation

$$\mathcal{L}f := f'' + \lambda|f'|^q - \frac{1}{2}\xi f' + \frac{q-2}{2(q-1)} f = 0. \tag{1.4}$$

2000 *Mathematics Subject Classification.* 34E05, 35K55.
Key words and phrases. KPZ equation, self-similar solutions.

We shall add to equation (1.4) the following initial data

$$f(0) = -1, \quad f'(0) = 0. \tag{1.5}$$

A more general initial condition $f(0) = f_0 < 0$ may be transformed by introducing a new function $f/|f_0|$.
Put

$$C = \left[\frac{1}{\lambda(q-1)} \left(\frac{q-1}{q} \right)^q \right]^{1/(q-1)}. \tag{1.6}$$

Our aim is to investigate the asymptotic behavior of solution of (1.4), (1.5) for $q > 2$. The main result of the paper is following.

Theorem 1.1. *Let $q > 2$ and $f(\xi)$ be a solution of problem (1.4), (1.5). Then*

$$\lim_{\xi \to \infty} \frac{f(\xi)}{\xi^{q/(q-1)}} = C, \tag{1.7}$$

where C is defined in (1.6).

The behavior of self-similar solutions of type $u(x,t) = t^\alpha g(xt^\beta)$ for (1.1) has been analyzed in [11].

2. The proof of Theorem 1.1

We start with simple result which is used later on.

Lemma 2.1. *Let $f(\xi)$ be a solution of problem* (1.4), (1.5) *defined for $\xi \in [0, \bar{\xi})$. Then*

$$f'(\xi) > 0 \quad \text{and} \quad f''(\xi) > 0 \text{ for } \xi \in (0, \bar{\xi}). \tag{2.1}$$

Proof. It is obvious, $f''(0) = (q-2)/[2(q-1)] > 0$. Therefore by continuity $f''(\xi) > 0$ and $f'(\xi) > 0$ in the right-hand side of some neighborhood of point $\xi = 0$. We suppose that there exists ξ_0 such that $0 < \xi_0 < \bar{\xi}$, $f''(\xi) > 0$ for $0 < \xi < \xi_0$ and $f''(\xi_0) = 0$. Then $f'(\xi) > 0$ for $0 < \xi \le \xi_0$ and $f'''(\xi_0) \le 0$. Equation (1.4) at ξ_0 subsequently leads to a contradiction. □

Now we shall obtain the upper bound for $f'(\xi)$.

Lemma 2.2. *There exists $\xi_0 > 0$ such that*

$$f'(\xi) < \left\{ \frac{\xi}{2\lambda} \right\}^{\frac{1}{q-1}} \text{ for } \xi \ge \xi_0. \tag{2.2}$$

Proof. Lemma 2.1 implies that $f(\xi) \to \infty$ as $\xi \to \bar{\xi}$, and, there exists unique point $\xi_0 \in (0, \bar{\xi})$ such that $f(\xi) < 0$ for $\xi \in (0, \xi_0)$ and $f(\xi) > 0$ for $\xi \in (\xi_0, \bar{\xi})$. Substituting $f'' > 0$ and $f \ge 0$ in (1.4) yields $f' < \{\xi/(2\lambda)\}^{1/(q-1)}$ for $\xi \in [\xi_0, \bar{\xi})$. Thus, $\bar{\xi} = \infty$ and (2.2) holds. □

Changing variables in (1.4)

$$f'(\xi) = \xi^{1/(q-1)} g(t), \quad \xi = \exp t, \tag{2.3}$$

we get new equation

$$g'' + \frac{3-q}{q-1} g' - \frac{q-2}{(q-1)^2} g = \left\{ \frac{1}{2} g' - \lambda (g^q)' + \frac{1}{q-1} g - \frac{\lambda q}{q-1} g^q \right\} \exp(2t). \tag{2.4}$$

By (2.1), (2.2) and (2.3) there hold

$$g(t) > 0 \quad \text{for any } t \in \mathbb{R}, \tag{2.5}$$

$$g(t) < \left\{\frac{1}{2\lambda}\right\}^{\frac{1}{q-1}} \quad \text{and} \quad g'(t) > -\frac{g}{q-1} \tag{2.6}$$

for sufficiently large values of t. Put

$$C_0 = \left\{\frac{1}{\lambda q}\right\}^{\frac{1}{q-1}}, \quad C_1 = \left\{\frac{1}{2\lambda q}\right\}^{\frac{1}{q-1}}. \tag{2.7}$$

It is obvious, $C_0 > C_1$. Now we shall establish the asymptotic behavior of $g(t)$ as $t \to +\infty$.

Lemma 2.3. *Assume that $g(t)$ is defined in* (2.3). *Then*

$$\lim_{t\to+\infty} g(t) = C_0. \tag{2.8}$$

Proof. From a careful inspection of equation (2.4) we conclude that a local maximum of $g(t)$ can happen only when $g(t) > C_0$.

At first we suppose that $g(t)$ does not tend to C_0 as $t \to +\infty$ and $g(t)$ is monotonic solution of (2.4). Then there exists $\bar{C} \neq C_0$ such that $\lim_{t\to\infty} g(t) = \bar{C}$. It is obvious, for any $\varepsilon > 0$ and some $A > 0$ there exists a sequence $\{t_k\}$ with the properties:

$$\lim_{k\to\infty} t_k = +\infty,\ |g''(t_k)| \le A,\ |g'(t_k)| \le \varepsilon.$$

Passing to the limit in (2.4) as $t = t_k \to +\infty$ and choosing ε in a suitable way we get that left-hand side is bounded, while right-hand side tends to infinity if $\bar{C} \neq 0$. Let $\bar{C} = 0$. Using (2.6) we conclude from (2.4) that

$$g'' + \frac{3-q}{q-1}g' \ge \frac{g}{3(q-1)}\exp(2t) \tag{2.9}$$

for large values of t. Then for large values of k (2.9) implies

$$g(t_k) \le \alpha \exp(-2t_k), \tag{2.10}$$

where positive constant α does not depend on k. Setting $\xi_k = \exp t_k$, from (2.3), (2.10) we get

$$f'(\xi_k) \le \alpha \xi_k^{(3-2q)/(q-1)},$$

that contradicts (2.1).

Now until the end of the proof we assume that $g(t)$ is not monotonic solution of (2.4). Suppose that $\liminf_{t\to\infty} g(t) < C_0$. Then there exist positive unbounded increasing sequences $\{s_k\}$ and $\{t_k\}$ such that $t_k > s_k$,

$$g'(t) \le 0,\ \ t \in [s_k, t_k], \tag{2.11}$$

and $g(s_k) = C_0,\ g(t_k) = C_\star$ for $k \in \mathbb{N}$, where $C_1 < C_\star < C_0$. Then

$$\frac{1}{2}g' - \lambda(g^q)' = -\lambda q(g^{q-1} - C_1^{q-1})g' \ge -\lambda q(C_\star^{q-1} - C_1^{q-1})g' \ge 0 \text{ on } [s_k, t_k]. \tag{2.12}$$

So, (2.4) and (2.12) imply that

$$g''(t) + \frac{3-q}{q-1}g'(t) \ge -\lambda q(C_\star^{q-1} - C_1^{q-1})g'(t)\exp(2s_k),\ \ t \in [s_k, t_k].$$

Hence, integrating with respect to t from s_k to t_k, we get

$$\left\{g'(t) + \frac{3-q}{q-1}g(t)\right\}\Big|_{s_k}^{t_k} \ge \lambda q(C_\star^{q-1} - C_1^{q-1})(C_0 - C_\star)\exp(2s_k).$$

This leads to a contradiction, since (2.5), (2.6) and (2.11) imply that the left-hand side of last inequality is bounded, while the right-hand side becomes unbounded as $k \to \infty$.

Let us prove that $\liminf_{t\to\infty} g(t) = C_0$. Indeed, otherwise there exist $\varepsilon > 0$ and sequence $\{\tau_k\}$ of points of local minima $g(t)$ with the properties: $\tau_k \to +\infty$ as $k \to +\infty$ and

$$g(\tau_k) \geq C_0 + \varepsilon. \tag{2.13}$$

Passing in (2.4) to the limit as $t = \tau_k \to +\infty$ we get a contradiction.

To end the proof we show that $\limsup_{t\to\infty} g(t) = C_0$. Otherwise $\limsup_{t\to\infty} g(t) > C_0$. Then there exist unbounded increasing sequences $\{s_k\}$ and $\{t_k\}$ such that $t_k > s_k > 2$,

$$g'(s_k) = 0,\ g'(t_k) = 0,\ g'(t) \geq 0 \text{ for } t \in [s_k, t_k],\ g(t_k) > C_0 + \delta,\ |g(s_k) - C_0| < \varepsilon, \tag{2.14}$$

where $k \in \mathbb{N},\ \delta > 0$,

$$\varepsilon = \min\left\{\delta/2, \frac{q-1}{4C_0}\delta^2, \left[1 - (7/8)^{\frac{1}{q-1}}\right] C_0\right\}. \tag{2.15}$$

Without loss of a generality we can suppose

$$C_0 - \varepsilon < g(s_k) < C_0 \tag{2.16}$$

or

$$C_0 \leq g(s_k) < C_0 + \varepsilon. \tag{2.17}$$

Let (2.16) be valid. If (2.17) is realized the arguments are similar and simpler. Denote by $\bar{t}_k \in (s_k, t_k)$ points such that

$$g(\bar{t}_k) = C_0. \tag{2.18}$$

Applying Hölder's inequality we derive

$$\int_{\bar{t}_k}^{t_k} g'\, d\tau \leq \left(\int_{\bar{t}_k}^{t_k} (g')^2 \exp(2\tau)\, d\tau\right)^{1/2} \left(\int_{\bar{t}_k}^{t_k} \exp(-2\tau)\, d\tau\right)^{1/2}$$

and therefore

$$\int_{\bar{t}_k}^{t_k} (g')^2 \exp(2\tau)\, d\tau \geq 2\delta^2 \exp(2\bar{t}_k). \tag{2.19}$$

We multiply (2.4) by $g'(t)$ and integrate after over $[s_k, t_k]$. Using (2.7), (2.14)–(2.16), (2.18) and (2.19) we obtain

$$\begin{aligned}
-\frac{q-2}{2(q-1)} g^2(t_k) &\leq \frac{q-3}{q-1}\int_{s_k}^{t_k} (g')^2\, d\tau + \int_{s_k}^{t_k} (g')^2 \left[\frac{1}{2} - \lambda q g^{q-1}\right] \exp(2\tau)\, d\tau \\
&+ \frac{\exp(2\bar{t}_k)}{q-1}\int_{s_k}^{\bar{t}_k} \left[\frac{1}{2}(g^2)' - \frac{\lambda q}{q+1}(g^{q+1})'\right] d\tau \\
&\leq -\frac{1}{4}\int_{\bar{t}_k}^{t_k} (g')^2 \exp(2\tau)\, d\tau + \frac{\exp(2\bar{t}_k)}{q-1}\left(\frac{g^2}{2} - \frac{\lambda q}{q+1} g^{q+1}\right)\Big|_{t=s_k}^{t=\bar{t}_k} \\
&\leq \left[-\frac{\delta^2}{2} + \frac{\varepsilon C_0}{q-1}\right] \exp(2\bar{t}_k) \leq -\frac{\delta^2}{4}\exp(2\bar{t}_k).
\end{aligned}$$

Passing to the limit as $k \to \infty$ we get a contradiction with (2.6). □

From Lemma 2.3 and the definition of $g(t)$ it follows (1.7). Theorem 1.1 is proved. □

Remark 2.4. Theorem 1.1 demonstrates the optimality of Theorem 2.3 in [10]. The arguments are the same as in Remark 4.6 of that paper.

Our next result shows that equation (1.4) with initial data

$$f(0) = f_0 > 0, \quad f'(0) = 0 \tag{2.20}$$

has no global solution.

Theorem 2.5. *Let $q > 2$ and $f(\xi)$ be a solution of problem (1.4), (2.20). Then there exists $\xi_\star$ such that $0 < \xi_\star < +\infty$ and $f(\xi) \to -\infty$ as $\xi \uparrow \xi_\star$.*

Proof. Suppose that problem (1.4), (2.20) has infinitely extendable to the right solution $f(\xi)$. Using arguments of Lemma 2.1 we show that $f'(\xi) < 0$ and $f''(\xi) < 0$ for any $\xi > 0$. From (1.4) we obtain

$$f'''(\xi) < -\lambda \left(|f'(\xi)|^q\right)'. \tag{2.21}$$

After integration (2.21) over $[0, \xi]$ we conclude that

$$f''(\xi) < -\lambda |f'(\xi)|^q. \tag{2.22}$$

Integrating (2.22) over $[\xi_1, \xi]$ $(0 < \xi_1 < \xi)$ we infer

$$\frac{1}{(q-1)|f'(\xi_1)|^{q-1}} > \lambda(\xi - \xi_1).$$

Passing to the limit as $\xi \to \infty$ we obtain a contradiction which proves the theorem. □

References

[1] L. Amour, M. Ben-Artzi; *Global existence and decay for viscous Hamilton-Jacobi equations*, Nonlinear Anal. **31** (1998), 621–628.

[2] S. Benachour, P. Laurencot; *Global solutions to viscous Hamilton-Jacobi equations with irregular initial data*, Comm. Partial Differential Equations **24** (1999), 1999–2021.

[3] S. Benachour, B. Roynette, P. Vallois; *Solutions fondamentales de $u_t - \frac{1}{2}u_{xx} = \pm|u_x|$*, Astérisque **236** (1996), 41–71.

[4] M. Ben-Artzi; *Global existence and decay for a nonlinear parabolic equation*, Nonlinear Anal. **19** (1992), 763–768.

[5] M. Ben-Artzi, J. Goodman, A. Levy; *Remarks on a nonlinear parabolic equation*, Trans. Amer. Math. Soc. **352** (2000), 731–751.

[6] M. Ben-Artzi, P. Souplet, F. Weissler; *Sur la non-existence et la non-unicité des solutions du problème de Cauchy pour une équation parabolique semi-linéaire*, C.R. Acad. Sci. Paris Sér. I Math. **329** (1999), 371–376.

[7] M. Ben-Artzi, P. Souplet, F. Weissler; *The local theory for viscous Hamilton-Jacobi equations in Lebesgue spaces*, J. Math. Pures Appl. **81** (2002), 343–378.

[8] I. Corwin; *The Kardar-Parisi-Zhang equation and universality class*, Random Matrices Theory Appl. **1** (2012) 1130001, 76 p.

[9] B. Gilding, M. Guedda, R. Kersner; *The Cauchy problem for $u_t = \Delta u + |\nabla u|^q$, J*, Math Anal. Appl. **284** (2003), 733–755.

[10] A. Gladkov, M. Guedda, R. Kersner; *A KPZ growth model with possibly unbounded data: Correctness and blow-up*, Nonlinear Anal. **68** (2008), 2079–2091.

[11] M. Guedda, R. Kersner; *Self-similar solutions to the generalized deterministic KPZ equation*, Nonlinear Diff. Equ. Appl. **10** (2003), 1–13.

[12] M. Kardar, G. Parisi, Y.-C. Zhang; *Dynamic scaling of growing interfaces*, Phys. Rev. Lett. **56** (1986), 889–892.

[13] J. Krug, H. Spohn; *Kinetic roughening of growing surfaces, in: Solids far from Equilibrium* (edited by C. Godrèche), Cambridge University Press, Cambridge, 1991, pp. 479–582.

Alexander Gladkov
Department of Mathematics and Mechanics, Belarusian State University, Nezavisimosti avenue,4, 220030, Minsk, Belarus
E-mail address: gladkoval@mail.ru